

Quantum acoustic bremsstrahlung of impurity atoms in a Bose–Einstein condensate

I.E.Mazets

Ioffe Physico-Technical Institute, 194021 St.Petersburg, Russia

We study the process of scattering of two impurity atoms accompanied by generation of an elementary excitation in a surrounding Bose–Einstein condensate. This process, unlike the phonon generation by a *single* impurity atom, has no velocity threshold and can be regarded as a quantum acoustic analog of a bremsstrahlung in quantum electrodynamics.

PACS numbers: 03.75.Fi, 05.30.Jp, 67.60.-g

Sixty years ago Landau proposed a phenomenologic microscopic description of the effect of superfluidity [1]. He considered a massive body moving in a medium with the elementary excitation energy spectrum $\epsilon(q)$, where $\mathbf{q}$ is the kinetic momentum of an elementary excitation. The energy conservation law avoids excitation generation and, therefore, dissipation of the kinetic energy of the moving body, if the velocity of the body is less than the critical one, $v_{cr} = \min_q[\epsilon(q)/q]$. In the case of the Bogoliubov tipe spectrum of elementary excitations in a dilute Bose–condensed gas [2], v_{cr} coincides with the speed of sound in such a degenerate quantum gas. It has been known, however, that the Landau criterion applies in its strict form only to microscopic objects moving through a Bose–Einstein condensate (BEC) [3]. If the size of an object exceeds the healing length of the BEC, the object must be regarded as a macroscopic one, and its motion can result in generation of vortex pairs in the BEC. Reduction, due to effects of the moving object size, of the critical velocity with respect to the value given by the Landau’s theory has been demonstrated in the MIT group experiments [4], where a laser beam has been used as a macroscopic object stirring the BEC. From the other hand, the MIT group also performed an experiment [5] on probing BEC superfluidity by microscopic objects, namely, by atoms transferred from the BEC to the untrapped hyperfine state by a Raman laser pulse. A good agreement of the results with the Landau criterion was demonstrated.

The probability of an elementary excitation generation by an impurity atom in a dilute BEC has been calculated by Timmermans and Côté [6].

The processes described above are not related to collisions between impurity particles in the presence of a BEC. In the present paper we try to fill this gap in the theory of ultracold gases. We consider a homogeneous BEC of neutral atoms of the first kind at zero temperature with a small admixture of atoms of the second kind (impurity atoms) traveling through the BEC at different velocities. The mass of an atom of the j th kind is denoted by m_j . The number density of atoms in the BEC is n_1 .

Let us consider two colliding impurity atoms. Before the collision their momenta are $\frac{1}{2}\mathbf{P} \pm \mathbf{p}$, respectively, where $\mathbf{P}$ is the center-of-mass momentum and $\mathbf{p}$ is the

momentum of their relative motion in the center-of-mass frame of reference. If the collision is accompanied by a generation of an elementary excitation with the momentum $\mathbf{q}$, in the final state the center-of-mass momentum is $\mathbf{P}' = \mathbf{P} - \mathbf{q}$, and the momentum of relative motion $\mathbf{p}'$ is less by absolute magnitude than p , unlike a case of elastic collision. The Bogoliubov spectrum for elementary excitations gives $\epsilon(q) = \sqrt{q^2/(2m_1)[q^2/(2m_1) + 2m_1c_s^2]}$, where c_s is the speed of sound in the BEC. The number density of impurity atoms is assumed to be small enough to neglect interaction with them in the expression for the energy spectrum of elementary excitations in the BEC. The energy conservation law

$$\frac{\mathbf{P}^2}{4m_2} + \frac{\mathbf{p}^2}{m_2} = \frac{(\mathbf{P} - \mathbf{q})^2}{4m_2} + \frac{\mathbf{p}'^2}{m_2} + \epsilon(q) \quad (1)$$

always admits solutions with $q \neq 0$ and the initial velocities of the colliding impurity atoms less (and even much less) than the critical velocity $v_{cr} = c_s$.

For the sake of simplicity, we shall consider hereafter the case of $\mathbf{P} = 0$. In this case Eq.(1) takes the form

$$\epsilon(q) + \frac{q^2}{4m_2} + \frac{p'^2 - p^2}{m_2} = 0. \quad (2)$$

It is obvious that even if $p \ll m_2c_s$, the generation of a phonon with $q \leq p^2/(m_2c_s)$ is possible.

In Fig. 1 we plot the five diagrams describing the process mentioned above. There are three different kinds of vertices corresponding to the different multipliers appearing in the transition matrix element $\mathcal{M}(\mathbf{p}, \mathbf{p}')$. The first one is the only kind of vertex appearing in the diagrams in Fig. 1 (a, b), where it is denoted by small filled circles. It corresponds to the multiplier $g_{12}\sqrt{n_1}(u_K - v_K)/\sqrt{V}$. Here V is the quantization volume, $u_K = \sqrt{\frac{K^2/(2m_1) + m_1c_s^2}{2\epsilon(K)} + \frac{1}{2}}$ and $v_K = \sqrt{\frac{K^2/(2m_1) + m_1c_s^2}{2\epsilon(K)} - \frac{1}{2}}$ are the Bogoliubov transformation coefficients [2], $\mathbf{K}$ is the kinetic momentum of an elementary excitation, $\mathbf{K} = \mathbf{Q}$ for a virtual phonon in the intermediate state and $\mathbf{K} = \mathbf{q}$ for the actual phonon in the final state. The coupling constant g_{12} describes interaction of atoms of the 1st kind with atoms of the 2nd kind and is introduced according to the common definition $g_{ij} = 2\pi\hbar^2(m_i + m_j)a_{ij}/(m_im_j)$, $i, j = 1, 2$, where a_{ij} is the scattering

length for s -wave scattering of an atom of the i th kind on an atom of the j th kind. Then in Fig.1 (c, d) we see vertices denoted by large filled circles and proportional to g_{22}/V . In Fig.1 (e), there is a vertex denoted by an open circle and proportional to $g_{11}\sqrt{n_1}[u_q(u_Q u_{Q'} - u_Q v_{Q'} - u_{Q'} v_Q) - v_q(v_Q v_{Q'} - u_Q v_{Q'} - u_{Q'} v_Q)]/\sqrt{V}$.




FIG. 1. The five diagrams describing the process of scattering of two impurity atoms (solid lines) accompanied by emission of a phonon (dashed line).

It is worth to note that the diagrams of Fig. 1 (a, b) are in certain sense analogous of the diagrams describing bremsstrahlung of charged particles in quantum electrodynamics [7]; impurity atoms and BEC elementary excitations in our case play role of charged massive particles and photons, respectively. However, in our case, the attraction potential between two impurity atoms emerging due to virtual phonon exchange is a short-range potential. For a pair impurity atoms moving with velocities below c_s this potential is just of the Yukawa type, and its Fourier transform is [8]

$$\mathcal{U}_{eff}(Q) = -\frac{g_{12}^2}{g_{11}} \frac{1}{1 + [Q/(2m_1 c_s)]^2}. \quad (3)$$

Thus, the diagrams of Fig. 1 (a, b) are similar to those of Fig. 2 (c, d) where the direct interaction between impurity atoms is short-range, too. To make the analogy with quantum electrodynamics closer, we refer to the bremsstrahlung process in a collision between a charged particle and a neutral atom (see, e.g., the recent paper by Korol and co-workers [9] and numerous references therein related to ordinary as well as to polarizational bremsstrahlung). The last diagram shown in Fig. 1 (e) is distinct from the previous ones. It has no analog among third-order processes in quantum electrodynamics (only in fifth order a similar process appears, if annihilation of two virtual photon and creation of one photon in a final state require creation of an electron-positron pair in an intermediate state). Therefore, it is reasonable to represent the transition matrix element as the sum

$$\mathcal{M}(\mathbf{p}, \mathbf{p}') = \mathcal{M}^{(1)}(\mathbf{p}, \mathbf{p}') + \mathcal{M}^{(2)}(\mathbf{p}, \mathbf{p}'), \quad (4)$$

where the first term in the right hand side corresponds to the four diagrams of Fig. 1 (a — d) together, and the

second term corresponds to the diagram shown in Fig. 1 (e).

We consider here the case of impurity atom velocities less than the speed of sound in the BEC. Under this condition elementary excitations both in the final and intermediate state belong to the phonon range of the Bogoliubov spectrum, and their magnitudes differ significantly: $Q \approx |\mathbf{p} - \mathbf{p}'| \gg q$. For the diagram of Fig. 1 (e) the momenta of the two virtual phonons are practically opposite, since their sum $\mathbf{Q} + \mathbf{Q}' = \mathbf{q}$ is relatively small. Therefore the effects of Bose statistics of phonons are not pronounced for the diagrams with two elementary excitations in the intermediate state [Fig. 1 (b, e)], because they occupy essentially different modes.

If the relative velocity of the colliding impurity atoms is small compared to the sound velocity then we have also $Q \ll m_2 c_s$ and, hence, can the processes contributing to $\mathcal{M}^{(1)}(\mathbf{p}, \mathbf{p}')$ within the simplified model of effective contact interaction. I.e., we describe them by the diagrams similar to those of Fig. 1 (c, d) where the vertex corresponding to the contact interaction of two impurity atoms is substituted by g_{eff}/V , where $g_{eff} = g_{22} - g_{12}^2/g_{11}$. It is also convenient to define the effective scattering length $a_{eff} = a_{22} - (1 + m_1/m_2)^2 m_2 a_{12}^2/(4m_1 a_{11})$, so that $g_{eff} = 4\pi\hbar^2 a_{eff}/m_2$.

Using the conditions and approximations discussed above, we calculate the matrix elements

$$\mathcal{M}^{(1)}(\mathbf{p}, \mathbf{p}') = -\frac{4g_{eff}g_{12}\sqrt{n_1}}{V^{3/2}c_s\sqrt{2m_1c_sq}}, \quad (5)$$

and

$$\mathcal{M}^{(2)}(\mathbf{p}, \mathbf{p}') = \frac{g_{12}^2 g_{11} n_1^{3/2}}{2V^{3/2}c_s^2\sqrt{2m_1c_sq}|\mathbf{p} - \mathbf{p}'|}. \quad (6)$$

For evaluation of an energetic denominator $(p^2 - p'^2)/m_2 + q^2/(4m_2)$ appearing in derivation of Eq.(5), the energy conservation law [Eq.(2)] was used. For the further calculations involving $\mathcal{M}^{(2)}(\mathbf{p}, \mathbf{p}')$, it is useful to recall the well-known expansions [10] $|\mathbf{p} \mp \mathbf{p}'|^{-1} = p^{-1} \sum_{l=0}^{\infty} (\pm p'/p)^l P_l(\cos \vartheta)$, where P_l 's are Legendre polynomials, ϑ is the angle between $\mathbf{p}$ and $\mathbf{p}'$, and $p' < p$. One can note that the matrix element given by Eq.(5) contributes only to s -wave scattering while the matrix element given by Eq.(6) contributes also to higher angular momentum scattering channels.

The total cross-section of the process of two impurity atoms collision accompanied by a phonon emission can be written, taking into account the effects of impurity atoms quantum statistics, as [11]

$$\sigma_{tot} = \frac{1}{j} \frac{2\pi}{\hbar} \int \frac{V d^3 \mathbf{p}'}{(2\pi\hbar)^3} \int \frac{V d^3 \mathbf{q}}{(2\pi\hbar)^3} \frac{1}{2} |\mathcal{M}(\mathbf{p}, \mathbf{p}') \pm \mathcal{M}(\mathbf{p}, -\mathbf{p}')|^2 \delta \left(\epsilon(q) + \frac{q^2}{4m_2} + \frac{p'^2 - p^2}{m_2} \right). \quad (7)$$

The probability flux density $j = 2p/(Vm_2)$ in the denominator is necessary for proper normalization of the cross-section. The upper sign in Eq.(7) applies if two colliding impurity atoms are identical bosons, and the lower sign applies if they are identical fermions.

After some tedious but straightforward calculations we get the total cross-section for a pair of colliding identical bosonic atoms:

$$\sigma_{tot}(p) = \sigma_s(p) + \sigma_{he}(p), \quad (8)$$

where the main term

$$\begin{aligned} \sigma_s(p) = & \frac{1}{30} a_{12}^2 \sqrt{4\pi n_1 a_{11}^3} \left(1 + \frac{m_2}{m_1}\right)^2 \left(\frac{p}{m_2 c_s}\right)^2 \\ & \times \left[\frac{a_{12}}{a_{11}} \left(1 + \frac{m_1}{m_2}\right) - 16 \frac{a_{eff}}{a_{11}} \frac{p}{m_2 c_s} \right]^2 \end{aligned} \quad (9)$$

describes s -wave scattering, and the relatively small correction

$$\begin{aligned} \sigma_{he}(p) = & \kappa_b a_{12}^2 \sqrt{4\pi n_1 a_{11}^3} \left(1 + \frac{m_2}{m_1}\right)^2 \left(1 + \frac{m_1}{m_2}\right)^2 \\ & \times \left(\frac{p}{m_2 c_s}\right)^2 \left(\frac{a_{12}}{a_{11}}\right)^2 \end{aligned} \quad (10)$$

describes scattering into partial waves with higher even angular momenta; $\kappa_b = \sum_{\nu=1}^{\infty} [2(4\nu+1)(4\nu+3)(4\nu+5)]^{-1} \approx 0.0023$.

In the case of two colliding identical fermionic atoms we get

$$\begin{aligned} \sigma_{tot}(p) = & \kappa_f a_{12}^2 \sqrt{4\pi n_1 a_{11}^3} \left(1 + \frac{m_2}{m_1}\right)^2 \left(1 + \frac{m_1}{m_2}\right)^2 \\ & \times \left(\frac{p}{m_2 c_s}\right)^2 \left(\frac{a_{12}}{a_{11}}\right)^2, \end{aligned} \quad (11)$$

where $\kappa_f = \sum_{\nu=0}^{\infty} [2(4\nu+3)(4\nu+5)(4\nu+7)]^{-1} \approx 0.0060$.

Since the process of collision of two impurity atoms accompanied by a phonon emission appears in higher orders of perturbation theory than a simple elastic collisions of two atoms (the cross-section of the latter process is of order of scattering length squared), the cross-section of quantum acoustic bremsstrahlung given by Eqs.(9, 10, 11) contains an extra small multiplier — the square root of the BEC diluteness parameter, $n_1 a_{11}^3 \ll 1$. An additional factor of smallness of this cross-section is the smallness of the ratio $p/(m_2 c_s)$ for slow collisions. However, it is possible to outline an experimentally feasible situation where the process of quantum acoustic bremsstrahlung cannot be neglected.

First of all, to suggest a realistic experiment, one needs not to forget about the effects of finite temperature. It is practically impossible to get in experiment the temperature lower than the chemical potential of a degenerate Bose-gas. At temperatures T of order of or larger than $m_1 c_s^2$, quantum acoustic bremsstrahlung never can be the

main process leading to thermal equilibration between the main component (atoms of the 1st kind) and the gas of impurity atoms. Certainly, the emission of phonons [6] by the supersonic fraction of thermal distribution of impurity atoms and the reciprocal process recently considered by Montina [12] dominate.

Thus we come to conclusion that a process involving bosonic stimulation has to be suggested.

Let us consider two counterpropagating plane matter waves of impurity atoms with atomic kinetic momenta equal to $\pm \mathbf{p}$, correspondingly. Each wave contains $N_2 \gg 1$ atoms, $n_2 = N_2/V$, and the total density of impurity atoms is $2n_2$. Additionally, we assume that in the BEC a sound wave with the momentum $\mathbf{q}$ is generated, and the phonon number in this sound wave is denoted N_{ph} . We assume that both N_2 and N_{ph} are so large that any effect arising from interaction with the above-condensate fraction at the given temperature is negligible compared to the processes described below. The momenta of the impurity atoms matter wave and the sound wave are chosen to be small, so that $p < m_2 c_s$ and $\epsilon(q) + q^2/(4m_2) < p^2/m_2$. Also generation of a phonon with the momentum $\mathbf{q}$ during the collisions of the two impurity atoms with momenta $\pm \mathbf{p}$ is allowed by the energy conservation law.

Since in the final state there are many phonons in the sound wave, the quantum acoustic bremsstrahlung process with phonon scattering to this particular mode is Bose-enhanced by a factor of $N_{ph} + 1 \approx N_{ph}$, and can prevail over other channels, provided that N_{ph} is large enough. It leads to amplification of the sound wave. However, there are two processes leading to the sound wave attenuation: (i) phonon scattering on impurity atoms, and (ii) the process reverse to quantum acoustic bremsstrahlung, namely, absorption of a phonon followed by an increase of kinetic energy of relative motion of a pair of impurity atoms.

The evolution of number of phonons in the sound wave is governed by the equation

$$\dot{N}_{ph} = \dot{N}_{ph}^{(col)} + \dot{N}_{ph}^{(qb)}, \quad (12)$$

where the first term in the right hand side corresponds to phonon scattering on impurity atoms and the second term corresponds to quantum acoustic bremsstrahlung and its reverse process. The first term can be written explicitly as follows:

$$\dot{N}_{ph}^{(col)} = -8\pi a_{12}^2 n_2 m_1^{-1} q N_{ph}. \quad (13)$$

This expression is exact if $m_2 \gg m_1$. However, even if m_1 and m_2 are comparable, Eq.(13) provides a reasonable approximation, provided that $p \ll m_2 c_s$.

Calculation of the rate of change of phonon number in the sound wave due to the processes involving a pair of impurity atoms results in the following expression:

$$\dot{N}_{ph}^{(qb)} = \frac{N_{ph} N_2^2 V m_2}{2\pi \hbar^2} \left[\frac{p'}{2} |\mathcal{M}(\mathbf{p}, \mathbf{p}') \pm \mathcal{M}(\mathbf{p}, -\mathbf{p}')|^2 \right]$$

$$\times \Theta \left(\frac{p^2 - p'^2}{m_2} - \epsilon(q) - \frac{q^2}{4m_2} \right) - \frac{p''}{2} |\mathcal{M}(\mathbf{p}'', \mathbf{p}) \pm \mathcal{M}(\mathbf{p}'', -\mathbf{p})|^2 \Big]. \quad (14)$$

Here $p' = [p^2 - m_2\epsilon(p) - q^2/4]^{1/2}$ is the momentum of a relative motion of an atomic pair after emission of a phonon, and $p'' = [p^2 + m_2\epsilon(p) + q^2/4]^{1/2}$ is the momentum after absorption of a phonon. For the sake of generality, we retain here the Heavyside's step function Θ , however, by assumption, its argument is positive. Bars over the expression denote averaging over the angle between kinetic momenta of relative motion of two impurity atoms before and after the inelastic collision. Plus or minus sign must be taken in the cases of bosonic or fermionic impurity atoms, respectively.

We write Eq.(14) more explicitly for the case of bosonic impurity atoms, since in this case the rate given by Eq.(14) is significantly larger than in the case of fermions. Taking into account that the isotropic scattering dominates for bosonic impurity atoms, we reduce Eq.(14) to

$$\dot{N}_{ph}^{(qb)} = \frac{\pi}{8} N_{ph} \frac{n_2^2}{n_1} a_{12}^2 c_s \frac{m_2 c_s}{p} \frac{m_1 c_s}{q} \left[\frac{p'}{p} \xi(p) \times \Theta \left(\frac{p^2}{m_2} - \epsilon(q) - \frac{q^2}{4m_2} \right) - \frac{p}{p''} \xi(p'') \right], \quad (15)$$

where $\xi(p) = \left(\frac{a_{12}}{a_{11}} - 16 \frac{a_{eff} p}{a_{11} m_2 c_s} \right)^2$. The net effect of the two processes contributing to Eq.(15) is always sound wave attenuation.

As an example, consider the case when $m_2 \approx m_1 \equiv m$, all the scattering lengths $a_{ij} \approx a_{12}$ (hence, $a_{eff} \ll a_{12}$), $p/(mc_s) \approx 0.3$, and $q/(mc_s) \approx 0.07$. If, e.g., we set $p = 0.3 mc_s$ and $q = 0.07 mc_s$, Eq.(12) gives

$$\dot{N}_{ph} = -8\pi n_2 a_{12}^2 c_s N_{ph} (0.07 + 0.21 n_2/n_1).$$

Thus increase of sound wave dissipation rate by few dozens percents seems experimentally feasible.

This work is supported by the Nederlandse Organisatie voor Wetenschappelijk Onderzoek, project NWO-047-009.010. One of the authors, I.E.M., also acknowledges the support from the Russian Foundation for Basic Research, grant No. 02-02-17686, and the Ministry of Education of Russia, grant No. UR.01.01.040. I.E.M. is grateful to Prof. D.A. Varshalovich and Prof. G.V. Shlyapnikov for helpful discussions.

- [3] D.R. Tilley and J. Tilley, *Superfluidity and Superconductivity* (IOP Publishing, NY, 1990).
- [4] C. Raman *et al.*, Phys. Rev. Lett. **83**, 2502 (1999); R.Onofrio *et al.*, Phys. Rev. Lett. **85**, 2228 (2000).
- [5] A.P. Chikkatur *et al.*, Phys. Rev. Lett. **85**, 483 (2000); see there also references to the previous works on probing superfluidity in liquid ^4He by admixed microscopic particles.
- [6] E. Timmermans and R. Côté, Phys. Rev. Lett. **80**, 3419 (1998).
- [7] V.B. Berestetskii, E.M. Lifshitz, and L.P. Pitaevskii, *Quantum Electrodynamics* (Butterworth Heinemann, Oxford, 1996).
- [8] M.J. Bijlsma, B.A. Heringa and H.T.C. Stoof, Phys. Rev. A **61**, 053601 (2000); H. Heiselberg *et al.*, Phys. Rev. Lett. **85**, 2418 (2000).
- [9] A.V. Korol *et al.*, J. Phys. B **34**, 1589 (2001).
- [10] D.A. Varshalovich, A.N. Moskalev, V.K. Khersonskii, *Quantum Theory of Angular Momentum* (World Scientific, Singapore, 1988).
- [11] E.M. Lifshitz and L.D. Landau, *Quantum Mechanics. Non-Relativistic Theory* (Butterworth Heinemann, Oxford, 1981).
- [12] A. Montina, e-print cond-mat/0204129.

-
- [1] L.D. Landau, J. Phys. (USSR) **5**, 71 (1941).
 - [2] N.N. Bogoliubov, J. Phys. (USSR) **11**, 23 (1947); for a recent review of the theory of Bose-condensed gases see F. Dalfovo *et al.*, Rev. Mod. Phys. **71**, 463 (1999).